

betaDiketonate-Iron(III) Complex: A Versatile Fluorine-19 MRI Signal Enhancement Agent.

Journal:	ACS Appl Bio Mater
Publication Year:	2019
Authors:	Chao Wang, Stephen R Adams, Hongyan Xu, Wenlian Zhu, Eric T Ahrens
PubMed link:	33981964
Funding Grants:	Molecular Imaging for Stem Cell Science and Clinical Application

Public Summary:

Fluorine-19 magnetic resonance imaging (MRI) has gained considerable momentum as a promising imaging modality for in vivo tracking of cellular therapies and as a diagnostic for inflammatory disease. To further the utility of this technique, we increase imaging probe sensitivity by merging paramagnetic metal chelates with aqueous perfluorocarbon (PFC) nanoemulsions. We prepared a highly fluorinated ferric tris(β -diketonate) chelate (MW = 1265.2 g/mol) at gram scale. This iron chelate is soluble in multiple PFC oils used for MRI and readily reduces the ^{19}F longitudinal relaxation time (T_1) to <100 ms with modest line broadening and displays superior properties for ^{19}F MRI applications. The sensitivity enhancement by Fe(III) laden PFC nanoemulsion was confirmed in MRI phantom studies, where reduced T_1 speeds data acquisition thereby increasing the ^{19}F image sensitivity per time via signal averaging. Additionally, ^{19}F relaxivity of nanoemulsions incorporating other metal ions, including Gd, Er, Ho, Dy, Mn, Cr, and Ni, were evaluated. High-moment lanthanide ions, such as Gd(III), display severe line broadening, but other ions [e.g., Ho(III)] induce pseudocontact chemical shifts (up to 0.5 ppm) of ^{19}F in nanoemulsion, which makes them potentially useful for multichromatic ^{19}F imaging. Formulated nanoemulsions have a shelf life >200 days. Free β -diketonate or its iron complex in formed PFC nanoemulsion did not induce cytotoxicity in intracellularly labeled macrophages. Overall, ferric tris(β -diketonate) chelate provides a scalable approach for boosting sensitivity of PFC-based ^{19}F MRI probes. More generally, it can functionalize PFC oil, whose chemical modification remains challenging.

Scientific Abstract:

Fluorine-19 magnetic resonance imaging (MRI) has gained considerable momentum as a promising imaging modality for in vivo tracking of cellular therapies and as a diagnostic for inflammatory disease. To further the utility of this technique, we increase imaging probe sensitivity by merging paramagnetic metal chelates with aqueous perfluorocarbon (PFC) nanoemulsions. We prepared a highly fluorinated ferric tris(beta-diketonate) chelate (MW = 1265.2 g/mol) at gram scale. This iron chelate is soluble in multiple PFC oils used for MRI and readily reduces the (^{19}F) longitudinal relaxation time (T_1) to <100 ms with modest line broadening and displays superior properties for (^{19}F) MRI applications. The sensitivity enhancement by Fe(III) laden PFC nanoemulsion was confirmed in MRI phantom studies, where reduced T_1 speeds data acquisition thereby increasing the (^{19}F) image sensitivity per time via signal averaging. Additionally, (^{19}F) relaxivity of nanoemulsions incorporating other metal ions, including Gd, Er, Ho, Dy, Mn, Cr, and Ni, were evaluated. High-moment lanthanide ions, such as Gd(III), display severe line broadening, but other ions [e.g., Ho(III)] induce pseudocontact chemical shifts (up to 0.5 ppm) of (^{19}F) in nanoemulsion, which makes them potentially useful for multichromatic (^{19}F) imaging. Formulated nanoemulsions have a shelf life >200 days. Free beta-diketonate or its iron complex in formed PFC nanoemulsion did not induce cytotoxicity in intracellularly labeled macrophages. Overall, ferric tris(beta-diketonate) chelate provides a scalable approach for boosting sensitivity of PFC-based (^{19}F) MRI probes. More generally, it can functionalize PFC oil, whose chemical modification remains challenging.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/betadiketonate-ironiii-complex-versatile-fluorine-19-mri-signal-enhancement>